
The Effect of Hpv on the Cervix

Akhmatova Gulrukh Rakhmatovna

Bukhara State Medical Institute, Uzbekistan

Abstract

The effect of HPV on the cellular composition of the cervix. Changes in the cervix in cancer and precancerous conditions are often asymptomatic and a routine examination will not always indicate pathology. Screening programs allow you to identify changes in a timely manner, determine the cause and prescribe treatment. Predisposition to cervical cancer is determined by genetic markers, but the contribution to the development of pathology is no more than 1% of all cases. External and internal factors have a great influence. Disorders of the hormonal background and tissue nutrition, the local state of immunity — determine the internal reason for which precancerous changes occur. Early active sexual life, frequent change of sexual partners and frequent childbirth, the use of hormone-containing contraceptives, the presence of STIs in the anamnesis are external risk factors for the development of cervical cancer. In 85% of cases, cervical cancer is associated with HPV.

Keywords: precancer, immunity, cervical cancer, HPV.

Introduction. HPV is a DNA-containing virus. The virus is related to epithelial cells. After entering the cells, the virus starts the synthesis of its own genetic material. Infected epithelial cells begin to actively divide. But due to violations, epithelial cells remain immature. The epithelial layer is not formed properly. Dysplasia occurs. In the cervix, in 90% of cases, the virus can be eliminated from the body due to the internal reserves of the woman's body. But prolonged presence of the virus in cells provokes cervical intraepithelial neoplasia (CIN) in 10% of cases. There are three histological classes of CIN: mild (CIN I), moderate (CIN II) and high (CIN III), with the transition of one to the other, the probability of cancer formation increases. Mild dysplasia (CIN I) in 90% of cases involute into normal tissue or remain unchanged, 10% progress to CIN II. CIN II is transformed into CIN III in 1 case out of 10. With HPV persistence for more than 3 years, cancer develops against the background of high-grade dysplasia. This situation is observed in half of women diagnosed with invasive cancer. There are about 200 types of HPV, but not every type is able to cause cervical lesions. HPV types 16 and 18 are usually detected in cervical intraepithelial neoplasia CIN III and invasive cancer, therefore it is called a highly oncogenic type virus. HPV types 6 and 11 are often associated with CIN I and CIN II and have low oncogenic potential.[1] Infection with the human papillomavirus (HPV) of a high degree of oncogenic risk is considered to be the main etiological factor in the pathogenesis of breast cancer. HPV refers to highly contagious infections with an incubation period of 3-4 weeks to 8 months. To date, more than 300 new papillomaviruses have been identified. Fourteen well-studied HPV genotypes (HPV 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66 and 68) are considered pathogenic or "high risk" genotypes and cause the development of cervical cancer. Chronic persistence of HPV types 16 and 18 causes cervical cancer in 70-75% of cases.

In the etiology of RSM, HPV infection is isolated. For the first time, papillomavirus infection was described in the I century BC by doctors of ancient Greece. In their opinion, genital warts were

transmitted sexually. In the 60s of the XIX century Melnick and A.D.Timofeevsky revealed the presence of virus-like particles in papillomas. In 1892, researcher Rigoni-Stern noticed that the cause of death in married women was often cervical cancer, but did not occur in virgins. Dunn researchers and Ogilvie in 1968 discovered viral particles in genital warts.

Results and Discussion. The presence of dysplasia is detected during a cytological examination of Papanicolaou (PAP smear) During the analysis, the features of the structure of cells, their morphology, location are studied, after which a conclusion is issued on the presence or absence of cell atypia.

To perform the analysis, the material must be obtained from three sites: the vaginal part of the cervix, the cervical canal and the transformation zone. The transformation zone is the transition zone of the multilayer flat epithelium of the cervix into the glandular epithelium of the cervical canal. Precancerous changes usually begin in the transformation zone.

The resulting material is transferred to a slide and stained, after which a cytologist examines the sample under a microscope. If the smear does not contain a cylindrical epithelium, then the sample is considered uninformative and is not subject to examination.

The smear sampling process is not standardized, so false negative results may be obtained. In order to avoid false answers, you need to follow the rules of sampling. The responsibility lies with the medical staff, the result depends on the qualifications of the person.

- Before the study, you cannot douche; inject medications and tampons into the vagina for 24 hours. During the day, it is necessary to exclude sexual intercourse;
- Smear sampling is performed prior to bimanual examination and extended colposcopy.

In this case, the sample will not contain foreign elements, and the laboratory will be able to issue a reliable conclusion. The result of cytological analysis

The cytological interpretation of the cellular composition should be presented to the attending physician in clinically relevant terms and definitions. There are several cytological classifications. The most common are the classification of Papanicolaou and Bethesda.

According to the classification of Pap smears are divided into 5 types:

- ✓ I — normal cervical epithelium of a healthy woman;
- ✓ II — elements of inflammation, mild atypia;
- ✓ III — dysplasia;
- ✓ IV — suspected cancer;
- ✓ V — cancer.

The Bethesda classification is international. It was created as a unified information transfer tool between cytologists and clinicians.

Conclusions. In accordance with WHO recommendations, the first cytological examination for Pap is performed at the age of 21 or three years after the onset of sexual activity. For the next three years in a row, it is recommended to undergo an examination once a year. If all three results are normal, then preventive examinations are carried out once every two to three years. After 40 years, the study must be taken at least once a year

References

1. Пантеева, Я. И., & Артамонов, Р. П. (2019). Роль вируса папилломы человека в развитии рака шейки матки. In *Бюллетень медицинских интернет-конференций* (Vol. 9, No. 9, pp. 374-375). Общество с ограниченной ответственностью «Наука и инновации».
2. Rakhmatovna, A. G. (2021). Efficiency of PDT in severe cervical dysplasia. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(3), 2566-2568.
3. DAVRONOVICH, D. R., AKHMATOVA, G. R., & SHOKIROV, B. S. (2020). Dynamics of the Immune Status of Women in the Treatment of Human Papilloma Virus (Hpv) of the Cervix. *JournalNX*, 6(06), 733-735.
4. Ахматова, Г. Р. (2022). ВЛИЯНИЕ РАЗНЫХ ФАКТОРОВ ПРИ ОБРАЗОВАНИЕ ЗЛОКАЧЕСТВЕННЫХ ОБРАЗОВАНИЙ ТИМУСА (ОБЗОР ЛИТЕРАТУР). *Scientific progress*, 3(3), 61-66.
5. Rakhmatovna, A. G. (2021). Efficiency of PDT in severe cervical dysplasia. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(3), 2566-2568.
6. DAVRONOVICH, D. R., AKHMATOVA, G. R., & SHOKIROV, B. S. (2020). Dynamics of the Immune Status of Women in the Treatment of Human Papilloma Virus (Hpv) of the Cervix. *JournalNX*, 6(06), 733-735.
7. Ахматова, Г. Р. (2022). ПРОГНОЗ ЦИТОКИНОВ ПРИ ЦЕРВИКАЛЬНЫХ ПОРАЖЕНИЙ ВИРУСА ПАПИЛОМЫ ЧЕЛОВЕКА (ВПЧ). *Scientific progress*, 3(4), 865-870.
8. Nurmurodovna, B. M. (2022). Morphological Changes in Various Tissues in Rats with Chronic I Nurmurodovna, B. M. (2022).
9. Morphological Changes in Blood Glucose and Liver Glycogen in Rats during Intoxication of Heavy Metal Salts. *Research Journal of Trauma and Disability Studies*, 1(9), 87-92ntoxication with Heavy Metal Salts. *Spanish Journal of Innovation and Integrity*, 7, 65-69.
10. Ахматова, Г. Р. (2022). ПРОГНОЗ ЦИТОКИНОВ ПРИ ЦЕРВИКАЛЬНЫХ ПОРАЖЕНИЙ ВИРУСА ПАПИЛОМЫ ЧЕЛОВЕКА (ВПЧ). *Scientific progress*, 3(4), 865-870.
11. Мухидова, Г. Х., & Тешаев, Ш. Ж. (2016). Морфометрические особенности параметров физического развития и антропометрических параметров верхней конечности 13 летних компьютернозависимых подростков мальчиков. *Problems of biology and medicine* (89) 2016г.стр, 52-56.
12. Мухидова, Г. Х. (2021). Сравнительная характеристика компьютерно-зависимых компьютерно-независимых подростков по антропометрическим показателям. *Тиббиётда янги кун*, 5, 37.
13. Khasanovna, M. G. (2021). Features of Computer Dependence Comparative Characteristics of Computer-Dependent and Computer-Independent Adolescents by Anthropometric Indicators. *International Journal of Innovative Analyses and Emerging Technology*, 1(6), 213-217.
14. Мухидова, Г. Х. (2021). Comparative Characteristics of morphometric parameters of physical development and anthropo extremities of healthy and computer.Research. *Jet Journal of Analysis and Inventions-RJAI*, 2(9), 14-17.
15. Мухидова, Г. Х. (2021). Феномен «компьютерной зависимости: особенности

- интернетзависимости у подростков». Eurasian journal of medical and natural sciences, 22-26.
16. Ilxomovich, N. A. (2022). CLINICAL AND EPIDEMIOLOGICAL FEATURES OF COVID-19 DISEASE IN CHILDREN. BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI, 2(9), 182-186.
 17. Nazarov, A. (2021). Healthy generation-the basis of a healthy family. Galaxy International Interdisciplinary Research Journal, 9(11), 409-413.
 18. Ilxomovich, N. A. (2022). CLINICAL AND EPIDEMIOLOGICAL FEATURES OF COVID-19 DISEASE IN CHILDREN. BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI, 2(9), 182-186.
 19. Ilkhomovna, N. F. (2022). Negative Impact of Seizures on Quality of Life. Miasto Przyszłości, 24, 120-122.
 20. Ilhomovna, F. N. (2022). LATE SEIZURES AND CONSEQUENCES OF EPILEPSY IN YOUNG CHILDREN. *Conferencea*, 219-223.
 21. Ilhomovna, F. N. (2022). RESPONSIBILITY OF PARENTS BEFORE THE OFFSPRING. *Conferencea*, 441-446.
 22. Nazarov, A. I. (2022). CONSEQUENCES OF SEIZURES AND EPILEPSY IN CHILDREN. *Web of Scientist: International Scientific Research Journal*, 3(02), 483-489.
 23. Nazarov, A. I. (2022). ATROF-MUHITNING INSON SALOMATLIGIGA TA'SIRI. *Scientific progress*, 3(1), 881-885.